

Full-Thickness Excision versus Shaving by Laparoscopy for Intestinal Deep Infiltrating Endometriosis: Rationale and Potential Treatment Options.

Autori: Laganà AS, Vitale SG, Trovato MA, Palmara VI, Rapisarda AM, Granese R, Sturlese E, De Dominicis R, Alecci S, Padula F, Chiofalo B, Grasso R, Cignini P, D'Amico P, Triolo O

Rivista: BioMed research international

DOI: 10.1155/2016/3617179

PMID: 27579309

ABSTRACT

Endometriosis is defined as the presence of endometrial mucosa (glands and stroma) abnormally implanted in locations other than the uterine cavity. Deep infiltrating endometriosis (DIE) is considered the most aggressive presentation of the disease, penetrating more than 5 mm in affected tissues, and it is reported in approximately 20% of all women with endometriosis. DIE can cause a complete distortion of the pelvic anatomy and it mainly involves uterosacral ligaments, bladder, rectovaginal septum, rectum, and rectosigmoid colon. This review describes the state of the art in laparoscopic approach for DIE with a special interest in intestinal involvement, according to recent literature findings. Our attention has been focused particularly on full-thickness excision versus shaving technique in deep endometriosis intestinal involvement. Particularly, the aim of this paper is clarifying from the clinical and methodological points of view the best surgical treatment of deep intestinal endometriosis, since there is no standard of care in the literature and in different surgical settings. Indeed, this review tries to suggest when it is advisable to manage the full-thickness excision or the shaving technique, also analyzing perioperative management, main complications, and surgical outcomes.